

From Ebola to other emerging infectious diseases: the case for strong risk assessment, mitigation and prevention



Yambuku Mission Hospital, DRC (Zaire), 1976



Yambuku Mission, DRC, 1976



Nurses, Yambuku Mission Hospital

Maternity, Yambuku Mission Hospital



Deceased health workers, Yambuku Mission Hospital, DRC, 1976



Sœur Beata
missionnaire à Yambuku
awei o Yambuku 19 sept. 1976



Sœur Myriam
missionnaire à Yambuku
awei o Kinshasa 30 sept. 1976

Soko motu akabi bomoi bwa yee mpo ya bandeko, nsuka ya bolingo wana (Yoh. 15.13).
Kristu alobi mpe asali bongo. Amipesi mpo ya bisu tee liwa iya kuruse.
Banyango ba bisu baneli: Sr. Beata, Sr. Myriam, Sr. Romana na Sr. Edmonda balandi, bamekoli Kristu.
Liloba liye iya yee likomi bomoi bwa bango. Ut'o bolenge bamipesi mobimba na Kristu na bosalisi
bakoni mpe batu ba mawa.
Eteko bomipesi boye boleki ndelu tee bokomi likama iya liwa, bakimi te, bamibendi nsima te kasi
batondisi fomeko la bolingani lokola 'te bobongoli mpo ya bango banso baneli libonza iya nsuka.
Sikawa o esengo ya yee, Kristu akotanga bango lisusu basaleli te, kasi bandeko.
Banyango ba bolingo, bandeko ba Kristu, bandeko ba bisu bopema na boboto mpe bosambela mpo
ya bisu tokoma lokola binu, bamekoli ba solo ba Kristu.



Sœur Romana
missionnaire à Yalosemba
awei o Yambuku 2 oct. 1976



Sœur Edmonda
missionnaire à Yambuku
awei o Kinshasa 14 oct. 1976



VROOM AANDENKEN AAN
Pater Germain LOOTENS
Missionaris van Scheut
geboren te St.-Kruis-Brugge op 30 oktober 1910,
priester gewijd op 18 augustus 1935,
naar Zaïre vertrokken op 14 augustus 1936,
overleden te Yambuku op 2 oktober 1976
als slachtoffer van een zware epidemie.

Ngaliema Hospital, Kinshasa, DRC




Filoform virus, first identified 1976, CDC (Atlanta) and Porton (UK)



Source: CDC

Animal market, near Yambuku, DRC




Patient record, outpatient department, Yambuku Hospital, DRC, August 1976

2348	Mago ba alima	♀	Bosanga	Lilongo	Helminthiase	27/8	31/8	4	-
2349	Alila Liwangu	♀	Bodaba	Shimbi	Bleno + Ankylo	27/8	31/8	4	-
2350	Mandungu Otundu	♀	yatuwama	Monzamboli	Ascariidose	27/8	31/8	4	J.
2351	Seambo Dombe	♀	yaongo	Biyowa	Bleno + Ankylostomiasis	27/8	31/8	4	-
2352	Ampidobolo Baka	♂	- 11 -	- 11 -	HI double	27/8	31/8	4	-
2353	Monzia Mokeka Gaga	♀	Boungulu	Lilongo	Ascariidose	27/8	31/8	4	-
2354	dingia Lidele	♀	yambawo	Monzamboli	HI D.	27/8	31/8	5	-
2355	Makilo Alita	♂	yandoupi	yandoupi *	epitaxin + dysentery	28/8	30/8	2	fur
2356	Koloupi Komlesa	♀	Kotaku	Lilongo	Bleno + Ascariidose	28/8	31/8	3	-
2357	Kanza K. Mubunzu	♂	yakolo	Monzamboli	contusion	30/8	31/8	1	-
2358	Batayo - Malike	♂	yaetoku	Moluwa	Anemie, + Ankylo	29/8	31/8	2	-
2359	Bunda Ozapi	♂	yambonzo	Biyowa	Malaria	30/8	31/8	1	-
2360	Opuwa Dosi	♀	yakai	Kwanza	Ankylost.	29/8	31/8	2	-
2361	Mapolola Mapula	♂	Celza jatoku	yandoupi	Ankylostomiasis	30/8	31/8	1	-
2362	Kebolo Ambena	♂	yamleka	- 11 -	HI Double	30/8	31/8	1	-
2363	Litinandunga Amba	♀	yalokila	Monzamboli	Observation	28/8	31/8	3	-
2364	Mondele Mohiwambi	♀	yapombi	- 11 -	Bleno + Ankylo	28/8	31/8	3	-
2365	Malene - Likonde	♀	yakoleka	yandoupi	Asp. thesion	28/8	31/8	3	-
2366	Eglogbo - Atalee	♂	yapbo	Monzamboli	Bronchite + Ascariid	29/8	31/8	2	-
2367	Ambena Gaya	♂	yandoupi	Monzamboli		30/8	31/8	1	-
2368	Boya - Makoma	♂	benzadi	yandoupi	Blessure plaie	30/8	31/8	1	-
2369	Aplapta A. Luaga	♀	yakombo	Monzamboli	Ankylostomiasis	30/8	31/8	1	-
2370	Likuja Soki	♀	yakombo	- 11 -	Helminthiase	30/8	31/8	1	-
2371	Zoda Mabambu	♀	- 11 -	- 11 -	Helminthiase	30/8	31/8	1	-
2372	Mangondo Mambo	♀	Bombanga	yandoupi	- Avortement	30/8	31/8	1	-

Hospital Implements, Yambuku, 1976



Ebola Haemorrhagic Fever by mode of transmission, Yambuku DRC, 1976



Risk assessment, Ebola haemorrhagic fever, 1976

- **Two highly lethal outbreaks simultaneously**
 - Zaire (Yambuku) 280/318
 - Sudan (Maridi) 151/284
- **Nosocomial transmission drove outbreaks into health workers and through them to community**
- **Animal reservoir suspected**
- **Unknown potential to reappear – one time emergence vs. periodic re-emergence**

Risk assessment, Ebola haemorrhagic fever, 1976

- **Two highly lethal outbreaks simultaneously**
 - Zaire (Yambuku) 280/318
 - Sudan (Maridi) 151/284
- **Nosocomial transmission drove outbreaks into health workers and through them to community**
- **Animal reservoir suspected**
- **Unknown potential to reappear – one time emergence vs. periodic re-emergence**

Mission Hospital, Tandala Zaire (DRC), 1977



1 clinical case/died

1 contact (sister) fit possible case definition/survived

1 historical probable clinical case/recovered, 1972




Ebola haemorrhagic fever surveillance, Zaire, 1981–1985: antibody in reported possible, probable and clinical cases

Case definition	1981 (<i>n</i> = 0)	1982 (<i>n</i> = 4)	1983 (<i>n</i> = 36)	1984 (<i>n</i> = 27)	1985 (<i>n</i> = 31)	1981–1985 (<i>n</i> = 98)
Possible	0	0	0	1	2	3
Clinical	0	1	4	2	4	11
Probable	0	2	5	0	0	7
Total	0	3	9	3	6	21

NOTE. *n* = no. of surveillance reports investigated.

Source: WHO

Ebola haemorrhagic fever surveillance, Zaire, 1981–1985: villages reporting probable, possible and clinical cases



Risk assessment, Ebola haemorrhagic fever, 1977

- **Two highly lethal outbreaks simultaneously**
 - Zaire (Yambuku) 280/318
 - Sudan (Maridi) 151/284
- **Nosocomial transmission can be prevented**
- **Animal reservoir suspected**
- **Periodic re-emergence occurs**


Kikwit General Hospital, Zaire, 1995



Nursing sisters, Kikwit General Hospital, Zaire, 1995



Ebola Haemorrhagic Fever by mode of transmission, Kikwit Zaire, 1995



Source: WHO/CDC

Ebola Haemorrhagic Fever, Mayibout Gabon, 1996



- **19 index cases: found and butchered freshly dead chimpanzee**
- **18 family members infected**
- **No nosocomial transmission**
- **21/37 (70%) fatal**

Tai Forest, Cote d'Ivoire, 1992



Chimpanzee die off, Tai Forest sociology research project area, 1992 - 1994



Risk assessment, Ebola haemorrhagic fever, 1994

- Periodic re-emergence occurs
- Highly lethal outbreaks occur periodically when health workers become infected
- Nosocomial transmission can be prevented
- Animal link to transmission confirmed
 - DRC (Yambuku and Tandala)
 - Cameroun



The search for a reservoir in nature, Ebola Haemorrhagic Fever, 1996



Source:: Emerging Infectious Diseases

The search for a reservoir in nature, Ebola Haemorrhagic Fever, 2001 - 2003

Journal home > Archive > Brief Communications > Abstract

Brief Communications

Nature **438**, 575-576 (1 December 2005) | doi:10.1038/438575a

Fruit bats as reservoirs of Ebola virus

Eric M. Leroy^{1,5}, Brice Kumulungui¹, Xavier Pourrut^{1,5}, Pierre Rouquet¹, Alexandre Hassanin², Philippe Yaba¹, André Délicat¹, Janusz T. Paweska³, Jean-Paul Gonzalez⁴ and Robert Swanepoel³

The first recorded human outbreak of Ebola virus was in 1976, but the wild reservoir of this virus is still unknown¹. Here we test for Ebola in more than a thousand small vertebrates that were collected during Ebola outbreaks in humans and great apes between 2001 and 2003 in Gabon and the Republic of the Congo. We find evidence of asymptomatic infection by Ebola virus in three species of fruit bat, indicating that these animals may be acting as a reservoir for this deadly virus. ▲ Top

1. Centre International de Recherches Médicales de Franceville, BP 769 Franceville, Gabon
2. Muséum National d'Histoire Naturelle, UMR 5202, Paris 75005, France
3. National Institute for Communicable Diseases, Special Pathogens Unit, Private Bag X4, Sandringham 2131, South Africa
4. Institut de Recherche pour le Développement, UR178, Mahidol University at Salaya, Nakhonpathom 73170, Thailand
5. Institut de Recherches pour le Développement, UR178, Centre International de Recherches Médicales de Franceville, BP 769 Franceville, Gabon

Correspondence to: Eric M. Leroy^{1,5} Email: eric.leroy@ird.fr

subscribe to
nature

ABSTRACT

Next ▶

+ Table of contents

Full text

Download PDF

Send to a friend

CrossRef lists 178 articles citing this article

Scopus lists 310 articles citing this article

Export citation


Scopus lists 310 articles citing this article

Rights and permissions

Risk assessment, Ebola haemorrhagic fever, 2002


- **Periodic re-emergence occurs**
- **Highly lethal outbreaks occur periodically when health workers become infected**
- **Nosocomial transmission can be prevented**
- **Animal link to transmission confirmed**
- **Bat probable reservoir in nature**

Selected Ebola outbreaks, 1976 - 2002



Source: WHO

Ebola outbreak, Guinea, December 2013 - present




Ebola outbreaks, West Africa 2014

Guinea	Sierra Leone	Liberia	Nigeria
506 cases	730 cases	599 cases	13 cases
373 deaths	315 deaths	323 deaths	2 deaths

Probable and suspected cases / confirmed deaths as of 9 August

Source: WHO



Initial Ebola economic impact, 2014

THE INDEPENDENT SATURDAY 23 AUGUST 2014

49

Sierra Leone, Liberia and Guinea had all been shaking off recent instable histories until Ebola decimated mining production, halted economic recovery and sent it back into reverse. PAULINE BAX, SILAS GBANDIA and ELISE ZOKER report from Freetown

Along came a virus and hauled three nations out of recovery


Sandi Sesay's boss promised him three months' pay when he told the driver to stop coming to work. The goal was to prevent any possible spread of the Ebola virus at the Sierra Leone mine that employs him.

Two weeks later, Mr Sesay, 29, has yet to see any money from Dawmus Construction, a contractor at London Mining's Marampa iron-ore deposit. "I take care of my mother, my sisters and my wife and three children," he said. "How am I going to cope?"

The prospects for both Mr Sesay and Sierra Leone were bright before the outbreak. The economy was set to grow 14 per cent, almost three times faster than the average in sub-Saharan Africa. In Liberia and Guinea, investment in iron ore was luring billions of dollars and fuelling growth.

Then the first case of Ebola appeared in December. Initially tagged as a short-term phenomenon, the disease now threatens to cripple three economies with a combined gross domestic product of about \$13bn (£8bn) - less than that of Afghanistan.

Commodity companies



tion of the three countries in the coming months.








The past few months mark the first time that the disease, identified in 1976 near the Ebola River in what is now the Democratic Republic of Congo, has killed anyone in West Africa. The virus lives naturally in fruit bats and other wild animals. Humans get it from the animal's secretions and pass it on to other humans through bodily fluids.

The outbreak is isolating the countries. Nigeria's Arik Air suspended flights to Liberia and Sierra Leone after a Liberian man travelled by plane to Lagos and infected at least eight others with the disease after he collapsed at the airport.

British Airways and Kenya Airways also halted routes to Liberia and Sierra Leone, while Gulf carrier Emirates scrapped flights to Guinea.

"It's not just that international flights are cancelled and movement of people is restricted because of the quarantine measures," said the political analyst Lansana Gberie. "There's also a disabling psychological atmos-

Breaches in species barrier since 1976

		Infection	Animal linked	Year 1 st reported
		Ebola virus	Bats	1976
		HIV-1	Primates	1981
		E. coli O157:H7	Cattle	1982
		Borrelia burgdorferi	Rodents	1982
		HIV-2	Primates	1986
		Hendra virus	Bats	1994
		BSE/vCJD	Cattle	1996
		Australian lyssavirus	Bats	1996
		Influenza A(H5N1)	Chickens	1997
		Nipah virus	Bats	1999
		SARS coronavirus	Palm civets	2003
		Influenza A(H1N1)	Swine	2009
		MERS coronavirus	? Camel	2012
		Influenza A(H7N9)	Chickens	2013

Nipah virus infection, Malaysia, 1998-1999



•Source: Chua KB, Journal of Clinical Virology, April 2003

Nipah virus outbreaks, humans, 1998 - 2008

Dates	Location	No. cases	No. deaths	CFR(%)
1998-1999	Malaysia;	265	105	40
1999	Singapore	11	1	9
2001	W. Bengal, India	66	45	68
2001	Bangladesh	13	9	69
2003	Bangladesh	12	8	67
2004	Bangladesh	29	22	76
	Bangladesh	36	27	75
2005	Bangladesh	12	11	92
2007	W. Bengal, India	5	5	100
2007	Bangladesh	15	8	54
2008	Bangladesh	11	6	54

Changing Nipah virus epidemiology: Bangladesh and India

- ✓ **Human-to-human transmission first suspected 2001, hospitalized patients, India**
- ✓ **Human to human transmission suspected again in 2003, 2005, and 2007, Bangladesh**
 - **cases could not be linked to domestic animal exposure, including pigs**
 - **index cases not identified: one potential exposure to bat guano in palm wine**

Assessing the risk/testing the hypothesis




Precautionary measures: community agreement to cover the collection containers



Community agriculture meeting




Swiss cheese events in epidemiology and public health



● James Reason: *BMJ* 2000;320:768-770

Shifting the paradigm from emergency response to prevention



Knowledge, attitude and practices study, 4 mining companies, DRC

Recognition of Impact of disease outbreak on mining companies:

ArcelorMittal, London Mining and African Minerals - postponed expansion plans and evacuated workers during current Ebola outbreak

Shares immediately fell in London trading

Clear recognition that a healthy community is a productive workforce

Community malaria control programmes reduce malaria-related work days lost by 94%; malaria-related spending at clinic by 76%;

Clear understanding of corporate social responsibility - “it is the right thing”

Must respond to NGO and other social pressures

KAP study, 4 mining companies, DRC

Clear understanding of potential barriers to improving current risk assessment/ mitigation/prevention

Costs because of demand for more services and replacement of government investment in public health

Corruption and lack of enforcement of regulations.

Clear understanding of facilitators to adopt risk assessment/mitigation and prevention strategies

Good practices in place in mining sites/camp to prevent, detect and control diseases.

Unambiguous company policies and enforcement (e.g. no bushmeat consumption in camp, provision of sufficient sources of protein in diet).

Ex Kulinda Afya – 11 Aug 2014

- One day desk top exercise: Katanga province DRC
- Objective: to raise awareness about emerging infections, their risk assessment, mitigation and prevention among senior field staff of mining companies and local government health officials.



Participants, Ex Kulinda Afya, 2014

- 25 participants: mining companies, provincial health authorities, animal health authorities, school of public health University of Lubumbashi
- Four groups with mining company, health and animal health representatives in each



Format of Ex Kulinda Afya exercise

Outbreak scenario (VHF) at mining site and town in a fictional African Country

Participants discussed and considered:

- Their initial response to outbreak of unknown disease
- Resources available to deal with an outbreak in community and in mining facilities
- Communication with and education of mine employees and surrounding areas
- How plans might be developed to mitigate the risk of future outbreaks



Conclusions of discussions during Ex Kulinda Afya

Internal risk mitigation procedures are effective in maintaining healthy workforce

External risk mitigation and preparedness procedures for outbreak alert and response are *ad hoc* and could be improved by:

- **Regional level:** increased cross working and coordination of public health activities between health representatives, the mining industries, and provincial representatives of non-governmental organisations.
- **Local level:** increased engagement and health education between the mining industries and the communities around the mining sites, with particular emphasis on risk assessment, mitigation, prevention and alert for zoonotic infections.
- **Sharing of financial, technical and logistical resources between mining industries and the provincial health authorities:**
 - equipment to assist in the isolation and quarantining of patients, and
 - access to laboratory testing

Potential role of mining companies in mitigation and prevention

Assessing risks from endemic and emerging infectious diseases in the communities and mining camps

Using available risk assessment resources, such as USAID toolkit, to optimise internal risk mitigation processes

Regular desktop scenarios/exercises to ensure external preparedness

Engaging (either individually or collectively) national and local governments as partners in infectious disease risk mitigation and prevention

Surveillance and alert networks in partnership with local communities

Health promotion/education/safe water/sanitation

Building trust

USAID Toolkit for assessment of internal risk management/mitigation processes

- To evaluate potential exposure points for diseases transmitted from animals and mitigate the risk of exposure
- Based on accepted best practice to address public health and environmental issues
- Adopting the practices could secure business continuity by securing the health of the workforce and neighbouring communities



There will always be a risk of emerging infections

There will always be a need to systematically consider emerging infectious diseases in business continuity plans

WHO: DRC Ebola outbreak in West Africa; cases jump

Filed Under: **Ebola**, **VHF**
Robert Roos | News Editor

Email Print & PDF

West Africa has seen a sharp increase in Ebola cases and deaths in the past 5 days.

The WHO said in a statement today. That compares with 24 suspected cases and 13 deaths reported on

